

Willkommen
Welcome
Bienvenue





PROBABILISTIC ASSESSMENT OF ENM FLOWS TO WASTE TREATMENT AND THE ENVIRONMENT

Véronique Adam, Bernd Nowack


Empa – Swiss Federal Laboratories for Materials Science and Technology
St. Gallen, Switzerland

Sustainable Nanotechnology Organization Conference – 07.11.2017







Based on Sun et al. 2014. Environmental Pollution




Based on Sun et al. 2014. Environmental Pollution




Based on Sun et al. 2014. Environmental Pollution







✓ Ag, CNT, CB, Cu, TiO₂, ZnO





✓ Ag, CNT, CB, Cu, TiO_2 , ZnO






✓ Ag, CNT, CB, Cu, TiO₂, ZnO





✓ Ag, CNT, CB, Cu, TiO₂, ZnO



Various sources:

- Reports
- Personal communications
- Papers
- Eurostat



Transfer coefficients

Various sources:

- Reports
- Personal communications
- Papers
- Eurostat



Transfer coefficients

France	Landfilling	Incineration	Recycling	Total
Packaging waste	16%	22%	62%	100%
Electronic waste	4%	6%	89%	100%

Various sources:

- Reports
- Personal communications
- Papers
- Eurostat



Transfer coefficients

Quality indicators:

Various sources:

- Reports
- Personal communications
- Papers
- Eurostat

Transfer coefficients

Quality indicators:

Geographical
correlation



Temporal
correlation



Reliability



Various sources:

- Reports
- Personal communications
- Papers
- Eurostat

Transfer coefficients

Quality indicators:

Geographical
correlation



Temporal
correlation



Reliability







Score

1

2

3







$$DQR = \frac{Geo + Temp + Rel + W_i \times 4}{i + 4}$$

W_i = weakest score

(EC, 2010)

i = number of indicators



$$DQR = \frac{Geo + Temp + Rel + W_i \times 4}{i + 4}$$

W_i = weakest score




(EC, 2010)

i = number of indicators



Data Quality Ratio	Coefficient of Variation
$1 \leq DQR < 1.5$	4.5%
$1.5 \leq DQR < 2.5$	13.75%
$2.5 \leq DQR \leq 3$	41.5%

(Laner et al., 2015)




$$DQR = \frac{Geo + Temp + Rel + W_i \times 4}{i + 4}$$

W_i = weakest score

(EC, 2010)

i = number of indicators



Data Quality Ratio	Coefficient of Variation
$1 \leq DQR < 1.5$	4.5%
$1.5 \leq DQR < 2.5$	13.75%
$2.5 \leq DQR \leq 3$	41.5%


(Laner et al., 2015)


Production mass

Production mass


	Comp1	Comp2	Comp3
Comp1	TRANSFER COEFFICIENTS		
Comp2			
Comp3			

	Comp1	Comp2	Comp3
Comp1	COEFFICIENTS OF VARIATION		
Comp2			
Comp3			






3'400 t nano-TiO₂ in solid waste in EU (mean)



MASS FLOWS AT NATIONAL SCALE



Summary

- Overall, high shares of ENMs enter sorting/recycling plants

Summary

- Overall, high shares of ENMs enter sorting/recycling plants
- Uncertainties are still quite high for some product applications and countries



Summary

- Overall, high shares of ENMs enter sorting/recycling plants
- Uncertainties are still quite high for some product applications and countries
- Adam & Nowack. 2017. *European country-specific probabilistic assessment of nanomaterial flows towards landfilling, recycling and incineration*. ES Nano 4:1961-1973

Summary

- Overall, high shares of ENMs enter sorting/recycling plants
- Uncertainties are still quite high for some product applications and countries
- Adam & Nowack. 2017. *European country-specific probabilistic assessment of nanomaterial flows towards landfilling, recycling and incineration*. ES Nano 4:1961-1973

Next steps


- Assess the mass flows out of waste treatment to the environment


Summary


- Overall, high shares of ENMs enter sorting/recycling plants
- Uncertainties are still quite high for some product applications and countries
- Adam & Nowack. 2017. *European country-specific probabilistic assessment of nanomaterial flows towards landfilling, recycling and incineration*. ES Nano 4:1961-1973

Next steps

- Assess the mass flows out of waste treatment to the environment
- Assess the **forms of releases**







5 forms

- **Dissolved**

5 forms

- **Dissolved**
- **Transformed**: subject to chemical reactions (e.g. sulfidation)

5 forms

- **Dissolved**
- **Transformed**: subject to chemical reactions (e.g. sulfidation)
- **Matrix-embedded**: in a solid matrix

5 forms

- **Dissolved**
- **Transformed**: subject to chemical reactions (e.g. sulfidation)
- **Matrix-embedded**: in a solid matrix
- **Nanoparticulate**: non-transformed, not embedded (free, aggregated and agglomerated)

5 forms


- **Dissolved**
- **Transformed**: subject to chemical reactions (e.g. sulfidation)
- **Matrix-embedded**: in a solid matrix
- **Nanoparticulate**: non-transformed, not embedded (free, aggregated and agglomerated)
- **Product-embedded**: contained within a product (solid waste)

5 forms



- **Dissolved**
- **Transformed**: subject to chemical reactions (e.g. sulfidation)
- **Matrix-embedded**: in a solid matrix
- **Nanoparticulate**: non-transformed, not embedded (free, aggregated and agglomerated)
- **Product-embedded**: contained within a product (solid waste)

5 forms




- **Dissolved**
- **Transformed**: subject to chemical reactions (e.g. sulfidation)
- **Matrix-embedded**: in a solid matrix
- **Nanoparticulate**: non-transformed, not embedded (free, aggregated and agglomerated)
- **Product-embedded**: contained within a product (solid waste)

Probabilistic assessment


















For each mass flow:















Probabilistic assessment


For each mass flow:



























Product category (nano-Ag)	Share of production (%)	Source
Electronics & Appliances	38	
Textiles	25	
Cosmetics	10	
Food	7	
Cleaning agents	6	
MedTech	4	
Plastics	3	
Paints	3	
Metals	2	
Glass & Ceramics	0.6	
Filter	0.3	
Diapers	0.2	



Product category (nano-Ag)	Share of production (%)	Source
Electronics & Appliances	38	
Textiles	25	
Cosmetics	10	
Food	7	
Cleaning agents	6	
MedTech	4	
Plastics	3	
Paints	3	
Metals	2	
Glass & Ceramics	0.6	
Filter	0.3	
Diapers	0.2	




Of mass released during use:





Product category (nano-Ag)	Share of production (%)	Source	Reliability
Electronics & Appliances	38		
Textiles	25		
Cosmetics	10		
Food	7		
Cleaning agents	6		
MedTech	4		
Plastics	3		
Paints	3		
Metals	2		
Glass & Ceramics	0.6		
Filter	0.3		
Diapers	0.2		

Of mass released during use:




Product category (nano-TiO ₂)	Share of production (%)	Source	Reliability
Cosmetics	59		
Paints	9		
Electronics & Appliances	7		
Cleaning agents	6		
Filters	6		
Plastics	4		
Coatings	4		
Glass & Ceramics	2		
Sporting goods	2		
Water treatment	0.7		
Food	0.4		
Textiles	0.3		
Spray	0.2		
Metals	0.1		
Cements	0.1		

Of mass released during use:




Production




33

Manuf. ▼




33

Use ▼




2.9

Export ▼




Nano-Ag (tons)







33




33




2.9




Nano-Ag (tons)




- Nanoparticulate
- Dissolved
- Transformed
- Matrix-embedded
- Product-embedded






TRANSFORMATIONS AT EUROPEAN SCALE



TRANSFORMATIONS AT EUROPEAN SCALE



Limitations

- Limited data, especially for nano-TiO₂
 - No size distribution
 - *Nanoparticulate* includes free, aggregated, agglomerated

Limitations

- Limited data, especially for nano-TiO₂
 - No size distribution
 - *Nanoparticulate* includes free, aggregated, agglomerated
- No outflows from recycling

Limitations

- Limited data, especially for nano-TiO₂
 - No size distribution
 - *Nanoparticulate* includes free, aggregated, agglomerated
- No outflows from recycling

Adam, Caballero-Guzman & Nowack. *Considering the forms of released engineered nanomaterials in material flow analysis*. Submitted to Nature Nanotechnology

Limitations

- Limited data, especially for nano-TiO₂
 - No size distribution
 - *Nanoparticulate* includes free, aggregated, agglomerated
- No outflows from recycling

Adam, Caballero-Guzman & Nowack. *Considering the forms of released engineered nanomaterials in material flow analysis*. Submitted to Nature Nanotechnology

Next steps

- Assessment of flows out of sorting and recycling plants
 - Masses and forms
 - Country-specific

Limitations

- Limited data, especially for nano-TiO₂
 - No size distribution
 - *Nanoparticulate* includes free, aggregated, agglomerated
- No outflows from recycling

Adam, Caballero-Guzman & Nowack. *Considering the forms of released engineered nanomaterials in material flow analysis*. Submitted to Nature Nanotechnology

Next steps



- Assessment of flows out of sorting and recycling plants
 - Masses and forms
 - Country-specific
- Dynamic probabilistic modeling




THANK YOU!



Empa

Materials Science and Technology



Score			
1	Country considered	≥ 2014	Reports, communications, papers
2	Local data	$2010 \leq x < 2014$	Eurostat
3	Different country	< 2010	Expert estimation